



Different Types of Cancer and their Characteristics

Zhuhai Li*

Department of Oncology, University of Chinese Academy of Sciences, Zhejiang, China

DESCRIPTION

Cancer is a complex disease that occurs when abnormal cells in the body begin to grow and divide uncontrollably. This can result in the formation of tumors or the spread of cancerous cells throughout the body, leading to a range of symptoms and potential health complications. There are many different types of cancer, each with its own unique characteristics, behaviors, and risk factors [1,2].

Breast cancer is the most common cancer among women worldwide, with approximately 2.3 million new cases diagnosed each year. It typically develops in the milk-producing ducts or lobules of the breast tissue. This can occur in both men and women, but it is more common in women. The symptoms of breast cancer can include a lump or thickening in the breast or underarm, nipple discharge or inversion, and skin changes on the breast. This type of cancer is highly treatable when detected early through regular mammography screenings, and treatment options include surgery, radiation therapy, chemotherapy, and hormone therapy. Risk factors include age, gender, family history of breast cancer, certain gene mutations, exposure to estrogen, and certain lifestyle factors such as obesity and alcohol consumption.

Prostate cancer is the most common cancer among men in the United States, with approximately 248,530 new cases diagnosed each year. It typically develops in the prostate gland, which is responsible for producing semen. This can occur in both younger and older men, but it is more common in men over the age of 65. The symptoms of this type of cancer can include difficulty urinating, weak or interrupted urine flow, and frequent urination. It is often slow-growing and may not cause any symptoms in its early stages. Treatment options for prostate cancer include surgery, radiation therapy, chemotherapy, and hormone therapy [3].

Risk factors include age, family history of prostate cancer, race, and certain lifestyle factors such as obesity and diet.

Lung cancer is the leading cause of cancer-related deaths in the world, with approximately 1.8 million deaths each year. It

typically develops in the cells that line the air passages of the lungs. Lung cancer can occur in both smokers and non-smokers, but it is more common in smokers. The symptoms of lung cancer can include coughing, chest pain, shortness of breath, and weight loss. This is often diagnosed at an advanced stage when it has already spread to other parts of the body, making it more difficult to treat. Treatment for lung cancer includes surgery, radiation therapy, chemotherapy, and targeted therapy [4-7]. Risk factors include smoking, exposure to secondhand smoke, exposure to certain chemicals and substances, and family history of lung cancer.

Colon cancer also known as colorectal cancer develops in the colon or rectum, which are part of the large intestine. It is the third most common cancer worldwide, with approximately 1.8 million new cases diagnosed each year. The symptoms of this cancer can include changes in bowel habits, rectal bleeding, abdominal pain, and unexplained weight loss [7-8]. Colon cancer is highly treatable when detected early through regular screening tests, such as colonoscopies. Treatment options for colon cancer include surgery, radiation therapy, chemotherapy, and targeted therapy. Risk factors for this type of cancer include age, family history of colon cancer, certain genetic conditions, and certain lifestyle factors such as diet, obesity, and physical inactivity. Leukemia is a type of cancer that develops in the blood and bone marrow, which are responsible for producing blood cells. There are several different types of leukemia, including acute lymphoblastic leukemia, acute myeloid leukemia [9, 10].

References

1. Hassanpour SH, Dehghani M. Review of cancer from perspective of molecular. *J Cancer Res Pract.* 2017;4(4):127-129.
2. Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-origin patterns dominate the molecular classification of 10,000 tumors from 33 types of cancer. *Cell.* 2018 ;173(2):291-304.
3. Friis S, Olsen JH, Thulstrup AM, Møller M, Linet M, Trichopoulos D, et al. Risk of liver and other types of cancer in patients with cirrhosis: A nationwide cohort study in Denmark. *Hepatol.*1998;28(4):921-925.

Correspondence to: Zhuhai Li, Department of Oncology, University of Chinese Academy of Sciences, Zhejiang, People's Republic of China, E-mail: danli@123.cn

Received: 26-Apr-2023, Manuscript No. BLM-23-21388; **Editor assigned:** 01-May-2023, Pre QC No. BLM-23-21388 (PQ); **Reviewed:** 15-May-2023, QC No. BLM-23-21388; **Revised:** 22-May-2023, Manuscript No. BLM-23-21388 (R); **Published:** 29-May-2023, DOI: 10.35248/0974-8369.23.15.570.

Citation: Li Z (2023) Different Types of Cancer and their Characteristics. *Bio Med.* 15:570.

Copyright: © 2023 Li Z. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

4. Jaouen G, Vessieres A, Top S. Ferrocifen type anti-cancer drugs. *Chem Soc Rev*. 2015;44(24):8802-8817.
5. Jaradat NA, Al-Ramahi R, Zaid AN, Ayesh OI, Eid AM. Ethnopharmacological survey of herbal remedies used for treatment of various types of cancer and their methods of preparations in the West Bank-Palestine. *BMC*. 2016;16(1):1-2.
6. Vigneri P, Frasca F, Sciacca L, Pandini G, Vigneri R. Diabetes and cancer. *Endocr. Relat. Cancer*. 2009;16(4):1103-1123.
7. Dano K, Behrendt N, Høyer-Hansen G, Johnsen M, Lund LR, Ploug M, et al. Plasminogen activation and cancer. *Thromb Haemost*. 2005;93(4):676-681.
8. Sak K. Cytotoxicity of dietary flavonoids on different human cancer types. *Pharmacogn Rev*. 2014 ;8(16):122.
9. Thomas L. On immunosurveillance in human cancer. *Yale J Biol Med*. 1982;55(4):329.
10. Nagy A, Munkacsy G, Gyorffy B. Pancancer survival analysis of cancer hallmark genes. *Sci Rep*. 2021;11(1):6047.